

2021 Moose Harvest Recommendation

to the
Vermont Fish and Wildlife Board



Vermont Fish and Wildlife Department
Agency of Natural Resources
1 National Life Drive, Davis 2
Montpelier, VT 05620-3208
802-828-1000

The Department's goal is to improve the health of moose in northeastern Vermont by reducing winter tick abundance and their impacts on moose health, survival, and birth rate. The Department recommends issuing a combination of either-sex and antlerless-only moose hunting permits in WMUs E1 and E2 to reduce the moose population and thereby reduce winter tick abundance.

The current number of moose in WMU E has been sufficient to sustain winter ticks at high levels that are negatively affecting moose health and survival. Winter ticks are a host-dependent parasite with moose being the primary host responsible for major fluctuations in winter tick densities. Therefore, reduction in moose density decreases the number of available hosts which in turn decreases the number of winter ticks on the landscape. Moose population reduction will be necessary to break the winter tick cycle and improve the health of moose in this region.

Failure to reduce the moose density will perpetuate the current, unhealthy state of the moose population in WMU E for decades and would be inconsistent with the Department's established objective of managing for a healthy moose population. Importantly, 65% of Vermont residents support maintaining a smaller moose population through hunting if it reduces the number of moose that die each year from winter ticks. Only 15% oppose this approach (Responsive Management 2019).

Although winter ticks can be found on moose throughout the northeast, they do not significantly impact moose populations across the more-peripheral parts of their range, including the rest of Vermont, due to lower moose densities that limit tick abundance.

Summary of Key Points

- The moose population is stable in most of Vermont and may be increasing in WMU E (E1 & E2).
- Moose density in WMU E remains above 1 moose/square mile.
 - No WMU outside of the Northeast Kingdom ever had a moose density of 1/mi².
 - Moose densities greater than 1/mi² support high numbers of winter ticks that negatively impact the health of moose.
 - Moose densities below 0.75/mi² support relatively few winter ticks that do not impact moose populations. This is the case in most of Vermont – winter ticks are present, but do not cause population level impacts.
- Results of moose research in WMU E indicate health of moose is very poor in that region.
 - Adult survival remains relatively good, but detrimental health impacts of winter ticks have caused birth rates to be very low.
 - About half of moose calves die each winter, primarily due to heavy winter tick loads.
- The Department recommends 100 moose hunting permits (60 either sex and 40 antlerless only) be allocated in WMU E to reduce moose numbers and thereby reduce the impacts of winter ticks on the health of moose and help maintain a sustainable moose population.
 - This would result in the harvest of 51-66 moose, or about 5% of the current estimated population in WMU E.
- No permits are recommended for the remaining 19 WMUs, which cover 93% of Vermont.

Goals

This recommendation aims to improve the health of moose in WMUs E1 and E2 by reducing the impact of winter ticks and to achieve moose population objectives established in the *2020-2030 Big Game Management Plan*.

Management Objectives

Moose population objectives for each WMU are established in Vermont's *2020-2030 Big Game Management Plan*. These objectives aim to maintain healthy regional moose populations at levels that are socially acceptable and ecologically sustainable.

In WMUs D2, E1, and E2, density objectives reflect the impact of winter ticks on the size and health of the region's moose population. Research has found reduced frequency of tick epizootics (where more than 50% of calves die from winter tick infestations) at moose densities below $1.06/\text{mi}^2$ and no tick epizootics at densities below $0.75/\text{mi}^2$ (Samuel 2007, Jones 2016). The Department will initially try to maintain moose densities at or below $1/\text{mi}^2$ to reduce winter tick abundance and the frequency of epizootics, and improve the health of the moose population. However, if tick impacts are not reduced, the moose density may need to be reduced to $0.75/\text{mi}^2$. Ultimately, the goal is to have healthy moose, with fewer calves dying each year from heavy winter tick loads and healthier cows with higher birth rates.

Moose density objectives throughout the rest of moose range in Vermont have been set at 0.5 moose/ mi^2 (Figure 1.). This lower objective reflects ecological limitations on moose densities in these regions due to limited young forest habitat, higher deer densities, and a warming climate. Moose densities in these WMUs have never reached $1/\text{mi}^2$.

Hunting thresholds have also been established for each WMU at 75% of the density objective. The Department will only consider hunting moose when densities exceed this threshold. This ensures that the other values of moose are maximized at these lower densities.




Figure 1. Moose density objectives (moose per square mile of moose habitat) established in Vermont's 2020-2030 Big Game Management Plan.

Population Status

Moose and Winter Ticks

Recent studies in Vermont, New Hampshire, and Maine have concluded that winter ticks are the primary cause of moose mortality across their core range in New England (Musante et al. 2007, 2010, Bergeron et al. 2013, Dunfey-Ball 2017, Jones et al. 2017, Ellingwood et al. 2019, Jones et al. 2019, DeBow 2020), with some moose hosting an astonishingly high number of ticks (>50,000/individual; Jones et al. 2019).

Core moose range (continuous red area in Figure 2) in New England extends from northeastern Vermont through northern New Hampshire and western and northern Maine. This part of the region has a colder climate with longer winters, low deer densities, large blocks of forest, and an abundance of young forest created by commercial timber management which allows it to sustain higher densities of moose than more peripheral parts of their range. Importantly, population-level effects of winter ticks have only been observed in the region's core moose range, where moose densities have been high enough to support large numbers of winter ticks.

Although winter ticks can be found on moose throughout the region, they are not impacting moose populations across the more-peripheral parts of their range in the northeast, including the rest of Vermont, due to lower moose densities which limit tick abundance. Moose numbers outside of the Northeast Kingdom have declined, but the main cause of that decline was not winter ticks. Rather, it was likely due to a combination of declining quantity of young forest, increased parasite loads (particularly brainworm linked to increasing deer densities), and fewer moose in core moose range to migrate out to these other regions.




Figure 2. Estimated probability of occurrence of moose in the New England region from Pearman-Gilman et al. 2020.

Vermont Research

During 2017–2019, 126 moose (36 adult cows and 90 calves) were fitted with GPS radio collars in WMU E to monitor survival and birth rates. Results of this research clearly showed that chronic, high winter tick loads have caused the health of moose in WMU E to be poor. Birth rates were low and overwinter calf survival was poor (49%; DeBow 2020). Although observed adult female survival remained relatively good, it was lower than expected for a population without major predators. Survival of breeding age females has significant influence on population trends in long-lived species like moose.

Much of the fieldwork associated with this research concluded in 2019; however, the Department continues to monitor survival and calf recruitment in the remaining collared cows. Additionally, University of Vermont researchers continue to analyze the large amounts of data collected during this

study to expand beyond the survival and health findings. Two additional research studies nearing completion are focused on understanding 1) How winter tick impacts on moose relate to habitat use and quality, and 2) How winter ticks affect moose genetic health and stress levels. Five scientific publications related to this research are currently in the final review stages, with more in production for 2022.

For more information about moose research in Vermont and New England, visit vtfishandwildlife.com.

Population Health

Many factors affect the health of individual moose and the overall population. These include diseases and parasites (e.g., winter ticks and brainworm), habitat quality, and environmental conditions. Ultimately, how fast a population grows and how resilient it is to additional sources of mortality is determined by how long individuals can be expected to live (i.e., the survival rate) and how many new individuals are added to the population each year (i.e., the birth rate).

In the early 2000s, moose were overabundant in WMU E. They were causing significant damage to forest regeneration and their physical condition was declining as habitat quality declined. The Department actively reduced the moose population in this area to bring it into balance with the habitat and to improve the health of moose. By 2011, the population had been reduced to a level the habitat could support; however, health measures did not improve (Figure 3).




Figure 3. Field-dressed body weights of harvested moose in Wildlife Management Unit E, 1993–2020.

Moose are not currently limited by habitat in the core part of their range, including WMU E (Dunfey-Ball 2017). There is enough available habitat and adequate forage to support the current population. However, habitat quality can influence the distribution of moose on the landscape (i.e., higher densities of moose in areas with the highest quality habitat), which can influence local winter tick abundance and impacts on moose health (Healy et al. 2019). Three publications from moose habitat research in Vermont examining how habitat quality influences moose distribution on the landscape and moose health will be forthcoming in 2021.

Based on comparable moose mortality research in New Hampshire and Maine during 2013–2017, it is possible that moose in WMU E experienced winter tick epizootics in 5 of 6 years from 2014–2019. Multiple years of heavy parasitism may have weakened the overall health of adults in the population and thus made them – and their calves – more susceptible to the effects of parasitism.

Tick Impacts in 2020 and 2021

The severity of annual tick infestations is not only dependent on moose density, but also on climate, including temperature, humidity, wind, and snow. Annual variation in climate conditions results in variation in winter tick loads on moose. As long as climate conditions periodically result in reduced winter tick infestations, moose numbers can continue to fluctuate at densities that perpetuate heavy tick loads and unhealthy moose for the foreseeable future.

Vermont did not have any collared moose calves in 2020. However, survival of collared moose calves in Maine was among the best observed since they began collaring moose in 2014. Anecdotal evidence (e.g., reports of dead moose, bloody beds, engorged ticks in snowmobile trails) suggests that tick impacts were lower in Vermont in 2020. That is supported by observed summer calf recruitment by collared cows, which was the highest since collaring began (4 years). Adult survival was also slightly better. It is likely that the long winter of 2018–2019 was a factor in reducing tick abundance during the winter of 2019–2020.

Winter tick counts on bull moose harvested in October 2020 were comparable to those observed in recent years (Figure 4). While this measure provides an indication of tick abundance on the landscape, final tick loads on individual moose will be largely determined by the length of the questing period. The questing period is typically ended by weather conditions (e.g., persistent snow or freezing conditions) that kill questing winter tick larvae. If the questing period ended early in 2020, tick loads will be moderate and the chance of an epizootic in 2021 will be low. However, if the questing period lasted into December, tick loads could be much heavier. It is not clear whether snow events in mid-October and early November in WMU E were sufficient to end the questing period.




Figure 4. Winter tick counts on bull moose harvested in Wildlife Management Unit E, 2013–2020.

Population Estimates

Regional moose densities in Vermont are estimated from moose sighting rates reported by deer hunters during the November rifle season. This approach was originally developed by the New Hampshire Fish and Game Department by relating sighting rates to moose densities determined by aerial surveys (Bontaites et al. 2000). Aerial surveys conducted in Vermont allowed the Department to modify this model to better fit Vermont sighting data. Sighting rates often vary from year to year due to factors other than the number of moose (e.g., weather conditions), so a 3-year rolling average is used to smooth out some of this variation.

Using this approach, the 2020 (2018–2020 rolling average) density estimates for WMUs E1 and E2 are 2.19 and 1.71 moose/mi², respectively, which are well above the upper density objectives established in the 2020-2030 *Big Game Management Plan* (1 moose/mi²; Table 1).

Moose densities in all other WMUs remain below established hunting thresholds (Table 1).

Table 1. Moose density estimates based on sighting rates by deer hunters and density objectives and hunting thresholds established in the 2020-2030 *Big Game Management Plan*, by WMU. Density estimates are based on average sighting rates during 2018–2020.

WMU	Habitat (mi ²)	Density (moose/mi ²)			Population Estimate	
		Objective	Hunting Threshold	Current Estimate	N	(80% CI)
A	35	n/a	n/a	0.03	1	(1–1)
B	420	n/a	n/a	0.06	25	(19–30)
C	351	0.5	0.38	0.32	112	(91–132)
D1	449	0.5	0.38	0.24	108	(89–128)
D2	346	0.75-1	0.56	0.33	116	(97–135)
E1	306	0.75-1	0.56	2.19	672	(611–732)
E2	326	0.75-1	0.56	1.71	558	(492–623)
F1	108	n/a	n/a	0.07	8	(5–11)
F2	158	n/a	n/a	0.04	6	(4–9)
G	363	0.5	0.38	0.06	23	(16–29)
H	466	0.5	0.38	0.34	159	(136–182)
I	407	0.5	0.38	0.13	52	(41–63)
J1	464	0.5	0.38	0.14	64	(51–77)
J2	633	0.5	0.38	0.25	158	(133–183)
K	359	n/a	n/a	0.04	14	(10–18)
L	346	0.5	0.38	0.25	85	(59–111)
M	424	0.5	0.38	0.27	114	(88–141)
N	275	n/a	n/a	0.04	11	(7–16)
O	478	n/a	n/a	0.03	16	(13–19)
P	447	0.5	0.38	0.12	55	(39–72)
Q	219	n/a	n/a	0.03	6	(4–7)
STATE	7380				2361	(2006–2719)

The Department has received interests for moose hunting from different foresters that have documented moose browse impacts to forest regeneration in different WMUs and are interested in alleviating these impacts to protect forest health. While some of these local areas could sustain a limited moose harvest, the moose population density in all WMUs except E1 and E2 remain below the hunting threshold.

The uneven distribution of functional moose habitat (and therefore moose) in parts of Vermont is a challenge for management. The Department will be reevaluating moose habitat mapping, taking advantage of recent research efforts (e.g., Pearman-Gilman et al. 2020, and forthcoming Blouin et al., currently in review) to better reflect the area of functional habitat in each WMU. This should allow for more meaningful estimates of moose density in WMUs with less homogeneous moose habitat.

Generally, it appears that moose numbers in WMU E have been relatively stable over the past 10 years (Figure 5). Given observed survival and recruitment rates from collared moose, it is unlikely that the moose population in WMU E is currently increasing as rapidly as sighting rates suggest. However, given limited moose harvests – and almost no cow harvest – since 2016, some increase in the population can reasonably be expected. Therefore, this harvest recommendation and all population projections herein are based on a more conservative projected population estimate of 1,000 moose (1.6 moose/mi²; approximately 80% of the 2020 estimate) in the fall of 2021.




Figure 5. Rolling 3-year average moose density estimates (solid line) and motor vehicle mortalities (dashed line) in WMU E during 2005–2020. Density estimates are based on moose sighting rates reported by deer hunters.

Harvest Recommendation

The results of the moose study clearly show that the current density of moose in WMU E has been sufficient to sustain winter ticks at high levels that are negatively affecting moose health and survival. Research has shown that winter tick abundance is directly related to moose population density. Reducing the density of moose decreases the number of available hosts which in turn decreases the number of winter ticks on the landscape. Moose population reduction will be necessary to break the winter tick cycle and improve the health of moose in this region.

Without management action to reduce the moose population, high tick loads will continue to impact the health of moose in WMU E for the next decade and beyond. The resulting chronic stress, low birth rates, and high calf mortality may prevent the population from growing. However, it will be less resilient to diseases, parasites, and environmental variation, which could cause the population to destabilize. Maintaining a healthy, stable, and sustainable moose population requires action to improve moose health.

Reducing winter tick numbers directly, either by treating moose or the landscape with some form of acaricide or fungal pathogen, is not currently a viable option. Research in this area is ongoing, but the realities of treating an entire landscape or a sufficient portion of the moose population make it unlikely that this will be a practical option in the near future.

The Department recommends harvesting at least 25 adult cow moose (5% of the cow population) in WMU E during the 2021 moose hunting seasons. The Department further recommends that this be accomplished through the issuance of 60 either-sex hunting permits and 40 antlerless-only hunting permits. Given historical success rates and sex-age composition of the harvest for each permit type, this allocation is expected to result in the harvest of approximately 58 moose (range: 51–66) with an expected breakdown of 28 bulls (range: 24–32), 26 cows (20–30), and 4 calves (3–6). Approximately 60% of permits are recommended to be allocated to WMU E1 due to higher moose densities in that WMU. Approximately 25% of either-sex permits are allocated to the archery season, based on the percentage of total applications that were for this season in recent years and the need to obtain sufficient biological data during the regular season. Allocations to the auction, special opportunity, and veterans are set by statute. Permit breakdown by season, type, WMU, and special allocation is provided in Table 2.

Table 2. Recommended 2021 moose hunting permit allocations by season, permit type, and WMU.

	E1	E2	Total
Regular Season¹			
Either-sex	24	15	39
Antlerless-only	24	16	40
Archery Season			
Either-sex	9	6	15
Auction²		choice	3
Special Opportunity³		choice	3
TOTAL			100

¹ Veteran permits are a priority draw for the first 5 regular season permits.

² Auction permits are either sex and winners have choice of season and WMU.

³ *Special Opportunity Permits are either sex and allow choice of season and WMU.*

Population Projections

Based on survival rates and calf recruitment observed from collared moose during 2017–2020, the moose population in WMU E would be expected to remain at its current density in the absence of any moose harvest (Figure 6), consistent with observed population trends over the past 10 years (Figure 5). If winter tick impacts are relatively severe each year (as observed during 2017–2019), tick-induced population declines, with no moose harvest, would take 12 years to reduce the population to 1 moose/mi². This presumably represents an unrealistic, worst-case scenario, and would be inconsistent with the observed population trend over the past decade. Further, detrimental effects on moose health will remain for several years after moose densities are reduced to levels that no longer support high tick loads. Even under this worst-case scenario, taking no management action will perpetuate the current, unhealthy state of the moose population in WMU E for many years and would be inconsistent with the Department’s established objective of managing for a healthy moose population. Importantly, 65% of Vermont residents support maintaining a smaller moose population through hunting if it reduces the number of moose that die each year from winter ticks. Only 15% oppose this approach (Responsive Management 2019).

Starting with a conservative projected population estimate of 1,000 moose in WMU E (E1 and E2 combined) in the fall of 2021, the harvest of 25 adult female moose annually is expected to reduce the population to 1 moose/mi² in 9 years, assuming tick impacts similar to the previous 4 years, and no improvement in birth rates or survival rates (Figure 6). If tick impacts are relatively severe each year, it would take 6 years at this permit allocation to reach 1 moose/mi². Conversely, if tick impacts are reduced, as in 2020, it would take 11 or more years at this allocation to reach 1 moose/mi².




Figure 6. Moose population projections in WMU E at varying annual cow harvests and winter tick impacts, based on a starting population of 1,000 and survival and birth rates from radio-marked moose. Projections assume consistent harvest each year and no improvement in survival or birth rates. Green

shaded area represents the potential range of variation due to varying winter tick impacts at the recommended harvest of 25 cows.

Given the poor health of the moose population and a clearly identified cause, action to address this issue is warranted. The number of permits allocated in 2020 was not sufficient to reduce the moose population in WMU E, even if 10 cows had been harvested, as expected, instead of 5. That conservative allocation was reflective of uncertainty around recent increases in population estimates, lower survival and birth rates observed from collared moose during the first 3 years of monitoring, and very low permit numbers in previous years.

The 2021 harvest recommendation is sufficient to reduce the moose population and thereby reduce winter tick impacts on moose in WMU E. However, it reduces the population slowly enough to allow for adjustments to the harvest, if necessary, even if the actual current density of moose is lower than 1.6 per square mile (1,000 moose). Ideally, moose health should be improved as quickly as possible. However, low survival and birth rates observed from Vermont moose, and broader, regional declines in moose populations justify a continued cautious approach at this time. Management of moose in WMU E and throughout Vermont must continue to be adaptive and respond to new information as it becomes available. If continued monitoring indicates that health, survival, and birth rates remain poor, and the moose population in WMU E remains above the objective, a more aggressive approach may be necessary to improve the health of the region's moose.

Literature Cited

- Bergeron, D. H., P. J. Pekins, and K. Rines. 2013. Temporal assessment of physical characteristics and reproductive status of moose in New Hampshire. *Alces* 49:39-48.
- Bontaites, K. M., K. A. Gustafson, and R. Makin. 2000. A Gasaway-type moose survey in New Hampshire using infrared thermal imagery: preliminary results. *Alces* 36:69-76
- DeBow, J. R. 2020. Effects Of Winter Ticks And Internal Parasites On Moose Survival And Fecundity In Vermont, USA. M. S. thesis. University of Vermont, Burlington, Vermont, USA.
- Dunfey-Ball, K. R. 2017. Moose density, habitat, and winter tick epizootics in a changing climate. M. S. thesis. University of New Hampshire, Durham, New Hampshire, USA.
- Ellingwood, D., P. J. Pekins, and H. Jones. 2019. Using Snow Urine Samples to Assess the Impact of Winter Ticks on Moose Calf Condition and Survival. *Alces*.
- Healy, C., P. J. Pekins, L. E. Kantar, R. G. Congalton, and S. Atallah. 2018. Selective habitat use by moose during critical periods in the winter tick life cycle. *Alces* 54:97-112
- Jones, H., P. J. Pekins, L. E. Kantar, M. O'Neil, and D. Ellingwood. 2017. Fecundity and summer calf survival of moose during 3 successive years of winter tick epizootics. *Alces* 53:85-98.
- Jones, H., P. Pekins, L. Kantar, I. Sidor, D. Ellingwood, A. Lichtenwalner, and M. O'Neal. 2019. Mortality assessment of moose (*Alces alces*) calves during successive years of winter tick (*Dermacentor albipictus*) epizootics in New Hampshire and Maine (USA). *Canadian Journal of Zoology* 97:22-30.
- Musante, A. R., P. J. Pekins, and D. L. Scarpitti. 2007. Metabolic impacts of winter tick infestations on calf moose. *Alces* 43:101-110.
- Musante, A. R., P. J. Pekins, and D. L. Scarpitti. 2010. Characteristics and dynamics of a regional moose *Alces alces* population in the northeastern United States. *Wildlife Biology* 16:185-204.
- Pearman-Gillman, S. B., J. E. Katz, R. M. Mickey, J. D. Murdoch, and T. M. Donovan. 2020. Predicting wildlife distribution patterns in New England USA with expert elicitation techniques. *Global Ecology and Conservation* 21.
- Responsive Management. 2019. Vermont residents' and hunters' attitudes toward big game hunting and management. Responsive Management Report, Harrisonburg, VA. 199pp.
- Samuel, W. M. 2007. Factors affecting epizootics of winter ticks and mortality of moose. *Alces* 43:39-48.